

1 **Differential expression of gonadotropin and estrogen receptors and oocyte cytology during**
2 **follicular maturation associated with egg viability in European eel (*Anguilla anguilla*)**

3 Filipa F.G. da Silva¹, Helge Tveiten², Gersende Maugars³, Anne-Gaëlle Lafont³, Sylvie Dufour³,
4 Josianne G. Støttrup¹, Elin Kjørsvik⁴, Jonna Tomkiewicz¹

5

6 ¹National Institute of Aquatic Resources, Technical University of Denmark, Kemitorvet, 2800 Kgs.
7 Lyngby, Denmark

8 ²Norwegian Institute of Fisheries and Food Research - Nofima AS, Muninbakken 9-13, Breivika, 9291
9 Tromsø, Norway

10 ³Museum National d'Histoire Naturelle, Sorbonne Universités, Research Unit BOREA, UPMC, CNRS,
11 IRD, UCN, UA, 7 rue Cuvier, CP 32, 75231 Paris, France

12 ⁴ Department of Biology, Norwegian University of Science and Technology, 7491 Trondheim, Norway

13

14 *Corresponding author:

15 Filipa F.G. da Silva

16 National Institute of Aquatic Resources, Technical University of Denmark, Kemitorvet, 2800 Kgs.
17 Lyngby, Denmark

18 Phone: +44 21125086

19 E-mail: fdsi@aqua.dtu.dk

20 **Abstract**

21 In captivity, oogenesis and ovarian follicle maturation in European eel can be induced experimentally
22 using hormonal therapy. The follicle's ability to respond effectively to the induction of maturation and
23 ovulation, resulting in viable eggs, depends on the oocyte stage at the time of induction. We
24 hypothesized that variation in the expression of key hormone receptors in the ovary and size of oocyte
25 lipid droplets are associated with changes in oocyte stage. Thus, we induced ovarian follicle maturation
26 using a priming dose of fish pituitary extract followed by an administration of 17α , 20β -dihydroxy-4-
27 pregnen-3-one (DHP). Females were then strip-spawned, the eggs were fertilized *in vitro*, incubated
28 and larval survival was recorded at 3 days post hatch (dph). The expression of gonadotropin receptors
29 (*fshr*, *lhcr1* and *lhcr2*) and estrogen receptors (*esr1*, *esr2a*, *esr2b*, *gpera* and *gperb*) was quantified
30 and the size of oocyte lipid droplets measured. Larval survival at 3 dph was used to differentiate high-
31 and low-quality egg batches. Results showed significantly higher abundance of *lhcr1* and *esr2a* at
32 priming for high-quality egg batches whereas *fshr* and *gperb* transcripts were significantly higher at
33 DHP injection for low-quality egg batches. Therefore, high levels of *lhcr1* and *esr2a* may be
34 important for attaining follicular maturational competence, while high *fshr* and *gperb* mRNA levels
35 may indicate inadequate maturational competence. Furthermore, lipid droplet size at DHP and in
36 ovulated eggs was significantly smaller in high-quality egg batches than in low-quality, which indicates
37 that droplet size may be a useful marker of follicular maturational stage.

38

39 **Key words:** estrogen receptors, gonadotropin receptors, lipid droplet, follicular maturation, egg quality

40 **1. Introduction**

41 The oceanic, reproductive stages of European eel (*Anguilla anguilla*) remain undiscovered and sexual
42 maturation neither occurs naturally in their continental habitats nor in captivity. This arrested
43 development results from a strong dopaminergic inhibition (Dufour et al., 1988; Vidal et al., 2004) and
44 a deficient pituitary gonadotropic function, i.e. both gonadotropin synthesis and release are low
45 (Dufour et al., 1983). Experimentally, gonadal development can be induced using hormonal treatments
46 based on fish pituitary extracts in females and human chorionic gonadotropin in males (Fontaine et al.
47 1964; Yamamoto and Yamauchi, 1974; Dufour et al., 1989; Ohta et al., 1996; Pedersen, 2003). In
48 females, such treatment leads to oocyte growth until maturation. Induction of follicular maturation and
49 ovulation generally requires an additional dose of pituitary extract and an injection of a maturation-
50 inducing steroid (MIS) (Yamauchi, 1990; Ohta et al., 1996; Pedersen, 2003) such as 17 α , 20 β -
51 dihydroxy-4-pregnen-3-one (DHP) in the case of the eel. Recent advances in eel assisted reproductive
52 technology have enabled the production of viable eggs and yolk-sac larvae (Butts et al., 2016; Sørensen
53 et al., 2016). However, resulting egg quality is variable and low fertilization rate and poor larval
54 survival are often observed. Variation in egg quality is partly related to the timing induction of ovarian
55 follicle maturation and ovulation, as previous studies have shown that hormonal treatment given too
56 early or too late in the reproductive cycle can be ineffective or inefficient (Palstra et al. 2005; Mylonas
57 et al., 2010; Unuma et al., 2011). Because hormones need to bind to receptors to exert their biological
58 function, differences in responsiveness to hormonal treatment could be due to differences in hormone
59 receptor expression.

60 In teleost fishes, as in other vertebrates, the two pituitary gonadotropin hormones follicle-stimulating
61 hormone (FSH) and luteinizing hormone (LH) play major roles in the regulation of oogenesis and

62 production of sex steroids (Pierce and Parsons, 1981). Gonadotropic function is mediated by specific
63 membrane receptors, the FSH receptor (FSHR) and the LH receptor (LHCGR), which corresponds to
64 the human LHCGR. These are mainly expressed in the somatic cells of the gonads (Rosenfeld et al.,
65 2007), though they can also be expressed in germ cells (Chauvigné et al., 2014). Until recently, it was
66 accepted that teleosts had a single FSHR and a single LHCGR encoded by *fshr* and *lhcr* genes,
67 respectively. However, in addition to *fshr*, Maugars and Dufour (2015) identified and characterized two
68 *lhcr* genes (*lhcr1* and *lhcr2*) in some fish species, including the European eel. European eel *lhcr1*
69 corresponds to the LHCGR previously characterized in the Japanese eel (*Anguilla japonica*) (Kazeto et
70 al., 2012). In comparison to the many studies on FSH and LH, knowledge about their receptors in
71 teleost species is still limited.

72 Sex steroids, such as estrogens, are also well-known for their role in reproductive function. In female
73 fish, estrogens (primarily estradiol-17 β , E2) are involved in the regulation of oogenesis and
74 vitellogenesis (Wallace, 1985) as well as in the feedback control of gonadotropin expression and
75 release (Zohar et al., 2010). Estrogens can diffuse through the cell membrane and bind to nuclear
76 estrogen receptors (ESRs). Two distinct subtypes of nuclear ESRs, *esr1* and *esr2*, have been cloned
77 from several mammalian and non-mammalian vertebrates (Mosselman et al. 1996; Hawkins et al.,
78 2000). In teleosts, including the European eel, two paralogs of *esr2* have been reported, *esr2a* (also
79 named *er β 2*) and *esr2b* (also named *er β 1*) (Hawkins et al., 2000; Ma et al., 2000; Menuet et al., 2002;
80 Nagler et al., 2007; Lafont et al., 2016). In addition, estrogens can also activate receptors on the cell
81 surface, initiating rapid and often non-genomic biological responses (Watson and Gametchu, 1999;
82 Falkenstein et al., 2000; Norman et al., 2004). The human G-protein coupled receptor 30 (GPER30 or
83 GPER) was shown to have the binding characteristics of an E2 membrane receptor (Revankar et al.,

84 2005; Thomas et al., 2005). Subsequent studies in zebrafish, Atlantic croaker (*Micropogonias*
85 *undulatus*) and common carp (*Cyprinus caprio*) have also shown that estrogens produced by follicle
86 cells inhibit or delay spontaneous follicular maturation via the activation of membrane receptors (Pang
87 et al., 2008; Pang and Thomas, 2009; Peyton and Thomas, 2011; Majumder et al., 2015). In the
88 European eel and some other teleosts, two paralogous *gper* genes (*gpera* and *gperb*) have been recently
89 identified (Lafont et al. 2016) but their roles have not yet been investigated.

90 During follicular maturation, cytological changes in the oocytes take place. This includes migration of
91 the germinal vesicle towards the oocyte periphery followed by breakdown of the nuclear envelope, and
92 meiotic resumption (Lubzens et al., 2017). In many teleosts, including the eel, another feature of oocyte
93 cytoplasmic maturation is the coalescence of lipid droplets to form one or a few large oil globules
94 (Kagawa, 2013). These cytological changes have been used as biomarkers for assessment of oocyte
95 maturational status in relation to assisted reproduction of eel (Palstra et al., 2005; Unuma et al., 2011).

96 The main objective of this study was to assess the expression of gonadotropin receptors (*fshr*, *lhcr1*
97 and *lhcr2*) and estrogen receptors (*esr1*, *esr2a*, *esr2b*, *gpera* and *gperb*) during induced maturation
98 and ovulation and investigate their relation with subsequent egg quality, ultimately estimated as
99 fertilization rate, hatching success and early larval survival. Finally, concomitant changes in oocyte
100 lipid droplet size were evaluated as potential cytological biomarker of follicular maturational status.

101 **2 Materials and Methods**

102 *2.1 Ethics statement*

103 All fish were handled in accordance with the European Union regulations concerning the protection of
104 experimental animals (Dir 86/609/EEC). Eel experimental breeding protocols were approved by the

105 Animal Experiments Inspectorate (AEI), Danish Ministry of Food, Agriculture and Fisheries (permit
106 number: 2010/561-1783). All efforts were made to minimize animal handling and stress.

107 2.2 *Experimental animals and rearing conditions*

108 Female silver eels (n=10; mean length and weight \pm SD were 72 ± 11 cm and 781 ± 393 g,
109 respectively) were caught from a freshwater lake (Vandet Sø) in northern Jutland (Denmark) and
110 transported to a research facility of the Technical University of Denmark located at Lyksvad Fish Farm
111 (Vamdrup, Denmark). Eels were randomly distributed into duplicate 300 L tanks equipped with a
112 recirculation system and gradually acclimated to artificial saltwater over a period of two weeks, i.e.
113 freshwater adjusted to 36 ppt salinity using Tropic Marin Sea Salt (Dr. Biener GmbH, Wartenberg,
114 Germany). Thirty male eels (body weight 106 ± 13 g; body length 38 ± 2 cm) reared on DAN-EX
115 2848 (BioMar A/S, Brande, Denmark) were obtained at a commercial eel farm (Stensgård Eel Farm
116 A/S, Randbøl, Denmark), transported to Lyksvad Fish Farm and kept in separate tanks under the same
117 conditions as the female eels. At the onset of hormonal treatment, the eels were anaesthetized
118 individually in an aqueous solution of benzocaine (ethyl p-aminobenzoate, 20 mg/L, Sigma-Aldrich,
119 Germany), tagged with a passive integrated transponder (pit-tag) in the abdominal muscle, and body
120 weight and length were measured. Throughout the experiment, all fish were maintained at ~36 ppt
121 salinity, ~20 °C, and a natural local daily photoperiod. No feed was provided during experiments since
122 eels in the migratory, silvering stage cease feeding (Lokman et al., 2003).

123 2.3 *Induction of gametogenesis and sampling*

124 Females received weekly intramuscular injection of salmon pituitary extract (SPE) at a constant dosage
125 of 18.75 mg/kg initial body weight to induce and sustain follicular development and vitellogenesis
126 (Kagawa et al., 2005; Tomkiewicz et al., 2012). The first injection was given concurrent with pit-

127 tagging and regular treatment lasted 16-20 weeks depending on the responsiveness of the females.
128 Pituitary extract was prepared using freeze-dried salmon pituitaries (Argent Chemical Laboratories,
129 Washington, USA) that were grinded, diluted in NaCl 0.9 g/L and centrifuged according to Ohta et al.
130 (1996, 1997). Supernatants were stored at -20 °C until use. Females were weighed at the weekly
131 injections to follow changes in body weight.

132 Individual treatment for follicular maturation and ovulation was initiated at first signs of the onset of
133 oocyte hydration, i.e. a body weight increase of 10 – 15 % compared to the initial weight and a soft
134 abdomen (Pedersen, 2003, 2004). To assess oocyte developmental stage, each female was
135 anaesthetized in an aqueous solution of benzocaine, and an ovarian biopsy (~0.2 ml) was obtained,
136 using a sterile disposable injection needle (16G x 1 ½”). The biopsy was taken at a standard location on
137 the left side of the body ~5 - 10 cm anterior to the genital pore, relative to female size, and the female
138 thereafter transferred to a separate tank under the same conditions for individual care. The biopsy was
139 inspected under the microscope and oocyte development graded on a scale from 1-7 according to
140 Palstra et al. (2005). Progression of oocyte maturation varied in time and homogeneity, so each female
141 was followed until the most developed oocytes exhibited characteristics close to stage 4, i.e., fully
142 transparent oocyte with nucleus at periphery (Fig. 1). At this stage, an additional SPE injection as
143 primer was given to females to sustain and boost follicular development (Pedersen, 2004; Kagawa et
144 al., 2005). To complete follicular maturation and induce ovulation, the eel maturation inducing steroid,
145 17 α , 20 β -dihydroxy-4-pregnen-3-one (DHP crystalline, Sigma-Aldrich Chemie, Steinheim, Germany)
146 was given ~24 h later at a dose of 2 mg/kg present body weight (Ohta et al., 1996). Prior to DHP
147 injection, a new biopsy (~0.2 ml) was obtained to evaluate the progression of oocyte development.
148 DHP was injected into the ovarian tissue under anaesthesia (Palstra et al. 2005). Female eels ovulated

149 12 to 15 hours after DHP injection and at that time the eggs were stripped by applying gentle pressure
150 along the abdomen of the fish. The volume of ovulated eggs collected was recorded.

151 Induction of spermatogenesis in the male eels started 4 weeks after the induction of oogenesis in
152 females, as males only need 7-9 weeks of treatment to reach spermiation (Tomkiewicz et al. 2011).
153 Each week, males received an intramuscular injection of human chorionic gonadotropin (hCG, Sigma
154 Aldrich Denmark, A/S) at dose of 1.5 IU hCG per kg initial body weight (Butts et al., 2014). An
155 additional hCG injection was given 12 h prior to milt collection in order to enhance sperm production.

156 2.4 *Sampling for gene expression and cytological analysis*

157 Sampling for each female comprised: 1) the ~0.2 ml ovarian biopsy obtained ~1h before the SPE
158 priming injection (SPE); 2) the ~0.2 ml biopsy obtained prior to DHP injection (DHP) and 3) a sample
159 of unfertilized eggs collected immediately after stripping (EGG). Each sample included digital images
160 of oocytes/eggs to measure lipid droplet diameter and a preserved sample (~0.1 ml) for gene expression
161 analysis. Prior to photography, the sampled oocytes/eggs were cleared in Serra's liquid
162 (ethanol:formalin:acetic at 6:4:1, diluted 20X in phosphate buffered saline, PBS, see Stoeckel (2000))
163 for staining of the germinal vesicle. The pictures were taken at 20X magnification using a digital
164 camera (Digital Sight DS-Fi1, Nikon Corporation, Japan) connected to an optical microscope (Eclipse
165 55i, Nikon Corporation, Japan) for measurement of lipid droplet diameter inside the oocytes. Samples
166 taken for analysis of gene expression were preserved in RNA-later (Ambion Inc., Huntingdon, UK),
167 refrigerated at 5 °C for 24 h and then frozen at -20 °C until RNA extraction.

168 2.5 *Fertilization rate, hatching success and larval survival*

169 In order to evaluate the egg viability, eggs were fertilized *in vitro*, and fertilization rate, hatching
170 success, and larval survival was determined. For each female, milt from 4-5 males was collected,
171 leaving three week intervals between individual male stripping. The spermatocrit was estimated
172 (pooled milt mixture according to Sørensen et al. (2013) and used to dilute milt in an immobilizing
173 diluent (Asturiano et al., 2004; Peñaranda et al., 2010). The diluted milt at a standardized concentration
174 was used for fertilization within 4 hours after collection (Butts et al., 2014).

175 Immediately after stripping, eggs were mixed with the milt solution and added natural seawater for
176 sperm activation (Butts et al. 2014). After 5 min gamete contact time, a sub-sample of eggs was
177 transferred to a 100 ml graduated cylinder glass to estimate percent floating eggs. The remaining eggs
178 were transferred to 10 L containers for separation of floating and sinking eggs. The eggs were kept at
179 20 °C and the fertilization rate and floating percent was accessed 3-5 hours post fertilization (HPF). To
180 estimate fertilization success, a sample of 100-150 floating eggs was photographed using an optical
181 microscope (Eclipse 55i, Nikon Corporation, Japan) at 20X magnification and a digital camera (Digital
182 Sight DS-Fi1, Nikon Corporation, Japan). Fertilized eggs were identified by the presence of blastomere
183 cleavage (minimum 4 cell stage), while those that had not reached the 4-cell stage were considered
184 unfertilized.

185 For estimation of hatching success, eggs (~200 eggs in triplicate, for each female) were collected from
186 the floating layer after 3-5 HPF and incubated at 20 °C in flasks (Nunc® 75 cm² flasks, non-treated
187 with ventilated caps, Thermo Scientific) (Sørensen et al. 2014). Each flask contained 250 ml of
188 seawater (36 ppt) ampicillin (50 mg/L) and rifampicin (50 mg/L). Flasks stayed undisturbed inside a
189 dark and closed incubator at 20 °C until the number of hatched larvae was counted at 55 HPF. To

190 estimate larval survival, ~300 g of eggs from the floating layer were incubated in a 60 L incubator with
191 natural seawater at ~20 °C. After hatch, when available ~3000 larvae in triplicate were stocked in 40 L
192 tanks of an aquaculture recirculation system containing natural seawater adjusted to 36 ppt salinity
193 using Tropic Marin Sea Salt and reared at 20 °C. Survival was measured as the percentage of still
194 living larvae at 3 days post hatch (dph).

195 For statistical analysis, egg batches were categorized into two groups: i) high-quality and ii) low-
196 quality based on larval survival. Hereby, survival beyond the first 3 dph (high mortality period) was
197 used to differentiate the groups (Table 2).

198 2.6 *Gene expression analysis*

199 RNA extraction and cDNA synthesis:

200 After Proteinase K treatment, total RNA was purified from ovarian biopsies and ovulated eggs by
201 homogenization of 30 mg of tissue in 700 µl TRIzol reagent (Invitrogen Life Technologies, Carlsbad,
202 CA, USA). TRIzol/chloroform separation was performed to remove DNA and proteins. The aqueous-
203 phase was transferred to a new tube and 500 µl of isopropanol were added for RNA precipitation.
204 Genomic DNA contamination was removed by treating the total RNA with the TURBO DNA-free kit
205 (Ambion). Total RNA quality and concentration was measured spectrophotometrically using
206 NanoDrop (Saveen Werner AB, Limhamn, Sweden).

207 RNA was then reverse transcribed into cDNA using Hight Capacity RNA-to-cDNA kit (AB) according
208 to manufacturer's protocol. In summary, reaction volumes of 20 µl contained 450 ng total RNA, 1X RT
209 buffer, 1X enzyme mix. The following cycling parameters were used: 37 °C for 60 min, 95 °C for 5
210 min and hold at 4 °C, using a 2720 Thermal cycler, AB.

211 Quantitative real-time PCR (qPCR):

212 The following gene specific primers were previously designed based on the nucleotide sequence of the
213 European eel: gonadotropin receptors, *fshr*, *lhgr1* and *lhgr2* (Maugars and Dufour, 2015); estrogen
214 receptors, *esr1*, *esr2a*, *esr2b*, *gpera* and *gperb* (Lafont et al., 2016). Specific primers for European eel
215 18S ribosomal RNA gene (*18S*) were designed in this study (Table 1), and their specificity and
216 efficiency was tested in qPCR.

217 The qPCRs were performed with a lightcycler (Roche, Ltd. Basel, Switzerland), using SYBR Green I.
218 Each reaction was prepared with 4 µl of diluted cDNA template, 2 µl PCR grade water, 2 µl of SYBR
219 Green master mix and 1 µl of each forward and reverse primers (0.5 pmol each at final concentration).

220 The following qPCR conditions were applied: polymerase activation step of 10 min at 95 °C, followed
221 by 51 cycles of 10 sec of denaturizing at 95 °C, 5 sec of annealing at 60 °C, 6 sec of elongation at 72
222 °C for *esr1*, *esr2a*, *esr2b*, *gpera* and *gperb* or 10 sec at 72 °C for *18S*, *lhgr1*, *lhgr2* and *fshr*. The
223 programs ended with a melting curve analysis by slowly increasing the temperature (0.1 °C/sec) from
224 65 °C to 95 °C, with a continuous registration of changes in fluorescent emission intensity. This last
225 step aimed at ensuring the presence of only one amplified product. Each qPCR run contained a non-
226 template control (cDNA was substituted by water) for each primer pairs to confirm that the qPCR mix
227 was not contaminated. Serial dilutions of a pool of ovary biopsies and ovulated egg cDNAs were used
228 as a standard curve for each gene. One dilution was included in each run as a calibrator. Normalization
229 of the qPCR data was performed using *18S* as a reference gene since it was stable through all sampling
230 points with a coefficient of variance at SPE, DHP and OV of 40%, 40% and 48%, respectively.

231 2.7 *Measurement of lipid droplet diameter*

232 Using the digital images of each sample (before SPE priming, before DHP injection and ovulated
233 eggs), 10 oocytes were randomly selected among those at the most advanced stage of development.
234 Here, ten of the largest lipid droplets were measured using the free software ImageJ (1.48d) and the
235 maximum five values averaged (Unuma et al., 2011). For each lipid droplet, the diameter was
236 calculated by the average of two diameter measurements. At the final stages of coalescence, only a few
237 droplets became larger while the others became smaller (Fig. 1), in these stages, the diameter was
238 based on the diameter of the largest droplets only (Unuma et al., 2011).

239 2.8 *Statistical analysis*

240 Statistical analysis was performed using R version 3.1.3 (R core team, 2015). Statistical differences in
241 gene expression and lipid droplet diameter between quality groups (high- and low-quality) in relation to
242 sampling time (before SPE, before DHP and after stripping) were evaluated using linear mixed-effects
243 (LME) models. The female ID was included in the models as within-subjects variable to account for
244 the repeated measurements taken on each female through the sampling points. Model assumptions of
245 normality and equal variance were checked using Q-Q plots and by observation of the residuals versus
246 fitted values plot. Square root and logarithmic transformations were applied to gene expression and
247 droplet diameter data whenever data deviated from a normal distribution. Linear regression analysis
248 was used to determine whether there were significant correlations between gene expression at each
249 sampling time and fertilization rate and hatching success. The expression of each gene was screened for
250 outliers using the Tukey's method. This method uses the interquartile range (IQR) to identify the
251 outliers range above and below the 1.5X IQR. A significance level (p) of 0.05 was applied in all tests.

252 3 Results

253 3.1 Egg production, fertilization and hatching success

254 The females studied produced on 327 ± 152 g eggs corresponding to 29-56 % of the initial body weight
255 (Table 2). The percent floating eggs varied from 25 to 100 %, while the fertilization rate ranged
256 between 15 to 99 % and the hatching rate between 0 to 80 %. Batch size was positively related to initial
257 female body weight ($R = 0.94$; $p < 0.001$) and length ($R = 0.95$; $p < 0.001$). Fertilization rate was not
258 correlated with female length ($R = -0.03$; $p = 0.930$) nor initial weight ($R = 0.09$; $p = 0.798$). Hatching
259 success ranged from 19 to 80 % and 0 to 5 % in the high- and low-quality groups, respectively (Table
260 2). There was also no significant correlation between hatching success and female length ($R = -0.03$; p
261 $= 0.927$) nor initial weight ($R = 0.05$; $p = 0.901$).

262 3.2 Expression of gonadotropin receptors

263 All three gonadotropin receptors (*fshr*, *lhcg1* and *lhcg2*) were detected in the ovarian biopsies with
264 genes differentially expressed across sampling points from the induction of oocyte maturation, SPE and
265 DHP, and EGG (Fig. 2). Overall, transcript levels of *fshr* did not vary during oocyte maturation, i.e.
266 between the ovarian biopsy taken before SPE and the biopsy taken before DHP ($p > 0.05$). However,
267 *fshr* levels dropped down below the level of detection of qRT-PCR detection in the eggs (at least 534
268 times less expressed than in the ovarian biopsies). Within sampling points, the expression levels of *fshr*
269 were similar between the high- and low-quality groups in the biopsies obtained prior to SPE priming (p
270 > 0.05 ; Fig. 2). However, before DHP, *fshr* transcript levels were significantly lower in the high-quality
271 group than in the low-quality group ($p < 0.001$). Present results also indicated a significant negative
272 correlation between *fshr* before DHP injection and hatching success ($R = -0.79$; $p = 0.012$; Table 3).

273 Overall, *lhcgr1* mRNA levels did not vary significantly ($p > 0.05$) during maturation (from SPE to
274 DHP injection). In contrast, transcript levels were around 150 times lower in ovulated eggs than in the
275 ovarian biopsies ($p < 0.001$). Moreover, in the ovarian biopsies collected before SPE priming, *lhcgr1*
276 expression was significantly higher in the high-quality group compared to the low-quality (p
277 = 0.002; Fig. 2). Present results also showed a significant positive correlation between *lhcgr1* at SPE
278 and hatching success ($R = 0.68$, $p = 0.032$; Table 3). There was no difference between the two groups
279 in the ovary samples taken before DHP injection and in ovulated eggs ($p > 0.05$).

280 Regardless of egg quality, *lhcgr2* transcript levels were similar between the ovarian biopsies taken
281 before SPE and DHP ($p > 0.05$) but significantly less expressed in the ovulated eggs (at least 5 times
282 less expressed than in the ovarian biopsies, $p < 0.001$). Within sampling points, transcript levels of
283 ovarian *lhcgr2* at SPE priming and DHP injection did not differ significantly between the high and low
284 quality groups ($p > 0.05$; Fig. 2). However, *lhcgr2* mRNA levels in unfertilized egg samples were
285 significantly higher in the high-quality than in the low-quality group ($p = 0.013$). There were no
286 significant correlations between *lhcgr2* transcript levels and hatching success or fertilization rate ($p >$
287 0.05; Table 3).

288 3.3 Expression of estrogen receptors

289 The nuclear receptor *esr1* was expressed in the ovarian samples obtained during oocyte maturation
290 without significant variation between the ovarian samples before SPE priming and DHP ($p > 0.05$; Fig.
291 3). After stripping, transcript levels were below the level of detection of qRT-PCR in unfertilized eggs
292 (at least 40 times less expressed than in the ovarian biopsies). Moreover, *esr1* expression was similar
293 between the high- and low-quality groups within the SPE and DHP sampling points ($p > 0.05$). There

294 were no significant correlations between *esr1* and hatching success or fertilization rate ($p > 0.05$; Table
295 3).

296 From the two nuclear *esr2* receptors, only *esr2a* was detected in the ovary during the induction of
297 oocyte maturation. Irrespective of hatching success, transcript levels of *esr2a* did not vary during
298 maturation treatment ($p > 0.05$), but the gene was much less expressed in unfertilized eggs (at least 67
299 times less expressed than in the ovarian biopsies, $p < 0.001$). In the biopsies taken before SPE priming,
300 transcript levels of *esr2a* were significantly higher in the high-quality group compared to the low-
301 quality group ($p = 0.020$; Fig. 3). In the ovarian biopsies taken before DHP and ovulated eggs, *esr2a*
302 expression levels were similar between groups ($p > 0.05$). We did not find any significant correlations
303 between *esr2a* transcript levels and hatching success or fertilization rate ($p > 0.05$; Table 3). Transcript
304 levels of *esr2b* were below the qRT-PCR detection threshold in all samples and could not be measured.

305 The two estradiol membrane receptors, *gpera* and *gperb*, were detectable and presented a different
306 expression pattern during induced oocyte maturation and after stripping (Fig. 3). Irrespective of egg
307 quality group, *gpera* transcript levels did not vary during maturation treatment ($p > 0.05$) but were
308 significantly lower in the eggs ($p < 0.001$). Expression of *gpera* did not differ significantly between
309 high- and low-quality in neither the ovarian biopsies before SPE and DHP injection, nor in ovulated
310 eggs ($p > 0.05$; Fig. 3). There were no significant correlations between *gpera* transcript levels and
311 hatching success or fertilization rate ($p > 0.05$; Table 3). Overall, there was no significant difference in
312 *gperb* mRNA expression levels during maturation ($p > 0.05$) while transcript levels of *gperb* dropped
313 to a level below the qRT-PCR detection threshold in ovulated eggs (at least 375 times less expressed
314 than in the ovarian biopsies). The expression levels of *gperb* before SPE administration did not differ
315 between the high-quality and low-quality group ($p > 0.05$) while levels were significantly lower in the

316 high-quality before the DHP administration ($p = 0.010$; Fig. 3). Present results did not show any
317 significant correlations between *gperb* levels and hatching success or fertilization rate ($P > 0.05$; Table
318 3).

319 3.4 Lipid-droplet related oocyte maturation status

320 Image analysis of the ovarian biopsies collected during the course of oocyte maturation induction
321 revealed 2-3 cohorts of oocytes of different sizes and stages of development (Fig. 4). Oocytes at the
322 most advanced stage of development were at the germinal vesicle migration stage, characterized by a
323 transparent cytoplasm and peripheral germinal vesicle visible before SPE and DHP injection (Fig. 1a
324 and b), similar to stage 4 and 5 according to the classification developed by Palstra et al. (2005),
325 respectively. No germinal vesicle could be observed in ovulated eggs after clearing in Serra's liquid
326 (Fig. 1c). As lipid droplets coalesced, their diameter increased significantly throughout oocyte
327 maturation and stripped eggs ($p < 0.001$). In the high-quality group, mean \pm SD lipid droplet diameter
328 was $89 \pm 23 \mu\text{m}$ at SPE priming, $136 \pm 14 \mu\text{m}$ at DHP injection and $156 \pm 15 \mu\text{m}$ at EGG. In the low-
329 quality group, lipid droplet diameter was $110 \pm 34 \mu\text{m}$ at SPE priming, $194 \pm 72 \mu\text{m}$ at DHP injection
330 and $248 \pm 74 \mu\text{m}$ at EGG. Overall, mean lipid droplet diameter was significantly higher and with wide
331 size variations in the low-quality than in the high-quality group ($p = 0.028$; Fig. 5), and mean size
332 increased significantly over time in both groups (from SPE to EGG, $p < 0.001$). Lipid droplet size in
333 the low-quality group also demonstrated an increasingly larger inter-female size variation over time,
334 compared to the high-quality group, where there were less lipid droplet size variation at all times. We
335 did not find a significant difference in droplet diameter between high- and low-quality at SPE priming
336 ($p > 0.05$) but at DHP and EGG, lipid droplet diameter was significantly higher in the low-quality
337 group ($p = 0.042$ and $p = 0.005$, respectively).

338 4 Discussion

339 4.1 Differential expression of gonadotropin receptors

340 Gonadotropin receptors mediate the biological effects exerted by gonadotropin hormones. In this study,
341 we analyzed for the first time the expression pattern of the *fshr* and the duplicate *lhgr* during the
342 induction of follicular maturation and ovulation of European eel using SPE treatment. The expression
343 of the two genes encoding distinct LHCGR (*lhgr1* and *lhgr2*) was also detected in the ovary of
344 immature eels (Maugars and Dufour, 2015). In the present case, both *lhgr1* and *lhgr2* were well
345 expressed in the ovary during induced maturation. Since the report of the coexistence of duplicated
346 *lhgr* in teleosts (Maugars and Dufour, 2015), their expressions have not yet been compared in any
347 other teleost species. In fishes, there are few studies about the expression of gonadotropin receptors
348 during follicular maturation and ovulation and none concerning gonadotropins and estradiol receptors
349 multiple paralogs. A previous study showed a high expression of ovarian *lhgr* orthologous to eel
350 *lhgr2*, in mature female Atlantic salmon (*Salmo salar*) caught during the upstream migration
351 (Maugars and Schmitz, 2006), which appears characteristic of the maturation stage prior to ovulation.
352 Moreover, we found that both genes were significantly less expressed in ovulated eggs. This agrees
353 with the location of LH binding sites found in the granulosa and thecal cells in European eel (Salmon et
354 al., 1988). Similarly, *lhgr* orthologous to eel *lhgr1* was found to be expressed in granulosa and
355 weakly in theca cells in chub mackerel (*Scomber japonicus*) (Nyuji et al., 2013) and Atlantic halibut
356 (*Hippoglossus hippoglossus*) (Kobayashi et al., 2008). Therefore, a drop in mRNA levels would be
357 expected in successfully ovulated eggs due to the loss of the follicle layers that remain inside the ovary
358 after the oocyte is extruded. Nevertheless, the drop in expression in ovulated eggs was more substantial
359 for *lhgr1* (150 less expressed) than for *lhgr2* (5 times less expressed), suggesting that *lhgr1* is
360 mostly expressed in follicular cells.

361 We hypothesized that differences in responsiveness to hormonal treatments during oocyte maturation
362 may be related to differences in hormone receptor expression, which consequently could affect
363 embryonic development. To test this, we calculated hatching success and larval survival as measures of
364 embryonic developmental competence (egg quality). Transcript levels of *lhgr2* during induction of
365 oocyte maturation were similar regardless of hatching success. However, *lhgr2* mRNA in ovulated
366 eggs was significantly higher in the high-quality group when compared to the low-quality group,
367 suggesting that *lhgr2* could have a role in embryonic development. Gonadotropin receptor transcripts
368 are also present in mouse oocyte and preimplantation embryo, with a potential beneficial role in oocyte
369 maturation and early embryonic development (Patsoula et al., 2001).

370 Previous studies have shown that gonadotropic treatment induces an increase in mRNA levels of *lh β* as
371 well as an increase in LH hormone in the pituitary in both European (Dufour et al., 1989; Schmitz et
372 al., 2005) and Japanese eel (Nagae et al., 1996; Saito et al., 2003; Jeng et al., 2007). This increase in
373 pituitary content of LH is likely necessary for the LH surge (in this case mimicked by the LH present in
374 the SPE) triggering ovulation. However, it is still uncertain if this observation is due to the effect of
375 gonadotropins alone, as both SPE and CPE treatments contain other components (neurotransmitters,
376 steroids) that can affect receptor expression. Our results showed that ovarian levels of *lhgr1* at the
377 time of SPE primer administration were both positively correlated with hatching success, and were
378 significantly higher in the high-quality group than in the low-quality group at SPE. This suggests that
379 females with higher hatching success had higher sensitivity to LH at the time SPE was administrated.
380 In contrast, the significantly lower levels of *lhgr1* expression in females with low-quality eggs could
381 have limited their response to the LH in the SPE primer. This is particularly important, in a first stage
382 of oocyte maturation, LH regulates the follicle's ability to produce maturation-inducing steroid (MIS)

383 and the oocyte's response to MIS (i.e. oocyte maturational competence); on a second stage the follicle
384 cells produce MIS (Patiño et al., 2001). Thus, failure to respond to LH at the first stage could
385 subsequently compromise the oocyte's ability to respond to the MIS (i.e. the DHP injection).
386 Additionally, LH signalling can stimulate other pathways than those leading to DHP synthesis. This has
387 been observed in zebrafish, where the expression of *ptgs2a* (a cox-2 gene) induced by LH was
388 necessary for ovulation to occur (Tang et al., 2017). Since oocyte maturation and ovulation may be
389 independently/differently regulated by the same ligand, the two processes may come out of synchrony
390 with each other, and this may influence the developmental capacity of the egg/embryo. In this case, the
391 mismatch between *lhcr1* expression and the administration of the SPE primer may have affected the
392 follicle's ability to respond to DHP in females in the low-quality group. Hence, a high expression of
393 *lhcr1* before SPE priming seems to be a good biomarker of high maturational competence.

394 Overall, *fshr* was well expressed in the mature ovary at all sampling times during the induction of
395 oocyte maturation. In Japanese eel, 17 weeks of treatment with pituitary homogenates also increased
396 ovarian levels of *fshr* (Jeng et al., 2007). However, the physiological relevance of high *fshr* expression
397 during ovarian maturation in teleosts is still unclear. In the case of the eel, gonadotropins contained in
398 exogenous pituitary homogenates (SPE in this case) could up-regulate the expression of both
399 gonadotropin receptors. Alternatively, in the case of multiple spawning, an ovulatory surge of FSH
400 could play an important role in the follicle recruitment for the next reproductive cycle (Prat et al., 1996;
401 Tyler et al., 1997; Sambroni et al., 2007). In this study, oocyte development presented a group-
402 synchronous pattern (Wallace and Selman, 1981), with 2-3 cohorts of oocytes of different size and
403 stage of development present in the ovarian biopsies taken during the induction of oocyte maturation
404 and ovulation. Thus, a high expression of *fshr* in the ovary during maturation could be related to the

405 regulatory role of FSH on the less developed oocyte batches also present in the biopsies. Alternatively,
406 it is also possible that FSHR has an important function on maturing oocytes (as occurs in mammals,
407 where FSH up-regulates LHCGR (Zelevnik et al., 1974). After ovulation, levels of *fshr* mRNA level
408 dropped below the level of detection, indicating that *fshr* is also mostly expressed in the follicle cells
409 surrounding the oocyte. This is in agreement with the observations by *in situ* hybridization of *fshr*
410 expression in granulosa and theca cells in vitellogenic follicles in salmon (Andersson et al., 2009).

411 In relation to hatching success, present results showed that transcript levels of *fshr* before the DHP
412 injection were negatively correlated with hatching success, and females with low quality eggs had
413 significantly higher transcript levels of *fshr*. This indicates that females in the low-quality group were
414 more sensitive to FSH at the time DHP was administrated. However, while *lhβ* mRNA levels tend to
415 increase, *fshβ* levels significantly decrease with gonadotropic treatment in both Japanese (Yoshiura et
416 al., 1999) and European eel females (Schmitz et al., 2005). Therefore, increased sensitivity to FSH at
417 the time of DHP injection does not appear to be determinant for successful oocyte maturation and
418 ovulation treatment in European eel. In contrast, increasing mRNA levels of *fshr* were associated to a
419 better competence of the oocyte to mature following pituitary hormone induction in rainbow trout
420 (*Oncorhynchus mykiss*) (Bobe et al., 2003). However, differences in hormone receptor expression
421 among species are likely influenced by differences in the reproductive strategies. For example, in
422 rainbow trout which is considered as total spawner (Mylonas and Zohar, 2007), *fshr* mRNA levels peak
423 at maturation and ovulation while *lhcgr* increased significantly later after ovulation (Sambroni et al.,
424 2007). In contrast, in zebrafish, which is a multiple batch spawner, *fshr* transcripts peaked at mid-
425 vitellogenesis and dropped at the end of vitellogenesis (Kwok et al., 2005). Overall, a combination of
426 low expression of ovarian *lhcgr1* before SPE priming and high expression of *fshr*, when DHP is

427 administrated, could be indicators of an ineffective response to oocyte maturation treatment resulting in
428 low egg quality.

429 4.2 Differential expression of estrogen receptors

430 In this study, we quantified gene expression of three nuclear (*esr1*, *esr2a*, *esr2b*) and two membrane
431 (*gpera* and *gperb*) estrogen receptors, during hormonal induction of oocyte maturation in European eel.
432 We found that transcripts of *esr1* were quite similar in the eel ovary throughout induced maturation,
433 with no significant differences between the two hatching groups. This suggests that *esr1* mRNA is not
434 a limiting factor for the development of maturational competence. Nevertheless, up-regulation of *esr1*
435 transcripts in the ovary of matured eels after ovulation (Lafont et al, 2015) suggests that this receptor
436 might play an important role during oocyte maturation. Regulation of *esr1* transcript levels may be
437 associated with an increase in circulatory levels of estradiol-17 β (E₂) during oocyte maturation stages
438 (da Silva et al., 2016). In particular, E₂ plasma concentration increases two to five times after SPE
439 priming in European eel (H. Tveiten, unpubl. results). Up-regulation of *esr1* was associated with an
440 increase of circulatory estrogens before ovulation also *in vitro* in eel hepatocytes (Lafont et al, 2015).
441 In contrast, *esr1* expression in the testis of male eels was markedly expressed in early stages of
442 spermatogenesis but significantly down-regulated in late stages of spermatogenesis (Morini et al.,
443 2017). This indicates a differential expression pattern of *esr1* between oogenesis and spermatogenesis
444 in eels. Differences in hepatic *esr1* expression between sexes have been observed in goldfish and are
445 likely related to differences in basal levels of E₂ between males and females (Nelson and Habibi, 2010).
446 In unfertilized eggs, *esr1* transcript levels were below the level of detection, suggesting that this
447 estrogen receptor is mostly expressed in the follicular cells.

448 Before SPE priming, *esr2a* transcripts were significantly more expressed in females with higher
449 hatching success which may indicate that this receptor has an important role during oocyte maturation.
450 In previous studies, the expression of *esr2a* in mature female eels after spawning was similar to
451 controls (Lafont et al., 2015) and in male eels, it was down-regulated during final stages of
452 spermatogenesis (Morini et al., 2017). Transcripts of *esr2a* were below the level of detection of the
453 qRT-PCR in ovulated eggs, suggesting that they are mainly expressed in follicular cells, as *esr1*.
454 Transcripts of *esr2b* were below the level of detection of the qRT-PCR in all samples, i.e. undetectable
455 during follicular maturation and ovulation. In male eels, *esr2b* transcripts in the testis were significantly
456 down-regulated throughout spermatogenesis (Morini et al., 2017). Thus, it is likely that *esr2b* does not
457 play a key role during the final stages of gametogenesis in European eel.

458 It is well known that in female fish GPER mediates the E₂-induced meiotic arrest of oocytes, e.g.
459 zebrafish, Atlantic croaker and common carp (Majumder et al., 2015; Pang and Thomas, 2009; Pang et
460 al., 2008; Peyton and Thomas, 2011). In these previous studies, only one GPER gene was
461 characterized, known now to be the orthologous to teleost *gpera*. We found that *gpera* transcripts were
462 expressed at similar levels in the ovary throughout the oocyte maturation (all stages), without
463 significant differences between the high-quality and low-quality group. Levels of *gpera* transcript
464 showed only a moderate decrease in ovulated eggs (not significant), indicating that *gpera* transcripts
465 are largely present in the eel oocyte itself, in contrast to all the other estradiol receptors. In contrast,
466 expression of *gperb* before DHP injection was significantly lower in females exhibiting higher egg
467 viability. Furthermore, *gperb* was the only estrogen receptor that was less expressed in females
468 exhibiting high quality eggs. This may suggest that *gperb* plays an important role in mediating the
469 inhibitory effect of estrogen during oocyte maturation also in European eel, where estrogen-dependent

470 oocyte growth is arrested at the first meiotic prophase by high levels of intracellular cyclic AMP
471 (cAMP) (Conti et al., 2002). In the pre-ovulatory phase, a LH surge causes a shift in the steroidogenic
472 pathway to the production of the MIS. When the MIS production (DHP injection in this case) is
473 sufficient to overcome the estrogen inhibitory effect, the binding of MIS to a G protein-coupled
474 progesterin membrane receptor triggers meiosis resumption and GVBD by causing a decrease in cAMP
475 concentrations (Jalabert and Finet 1986; Finet et al., 1988). Moreover, in Japanese eel, increased E₂ in
476 the late stages of oocyte development inhibits the production of DHP (Ijiri et al., 1995). Thus, a lower
477 expression of *gperb* in the HEQ group suggests a lower sensitivity to estrogen and subsequently a
478 lower estrogen inhibition, which may have enhanced the effect of DHP. In male eels, both *gpera* and
479 *gperb* transcripts significantly increased during spermatogenesis suggesting that GPERs may be
480 involved in final sperm maturation (Morini et al., 2017). Overall, our results raise the possibility that an
481 E₂ pathway, possibly regulated by *esr2a* and *gperb*, is involved in the control of oocyte maturation also
482 in European eel. However, we did not find any significant correlation between the expression of these
483 two genes and other egg quality indicators such as fertilization and hatching success. Thus, evidence is
484 weak and further investigation is necessary to clarify their role during oocyte maturation and
485 subsequent egg quality.

486 4.3 Cytological indicators of oocyte maturation status

487 An early index of maturational acquisition is the onset of germinal vesicle migration, which is
488 accompanied by the rise in LH at the end of follicle growth (Lubzens et al., 2010). In this study, a
489 peripheral germinal vesicle in the most advanced oocytes was used as an indicator that females had
490 progressed into initial stages of oocyte maturation before SPE priming. The appearance of oil droplets
491 was used as a marker of the maturation progress to initiate SPE priming or DHP injection (Palstra et al.

492 2005; Unuma et al., 2011). Here, the coalescence of the oil droplets reflects the hormonal processes
493 during final maturation resulting in decreasing numbers and increasing size of the lipid droplets.
494 Interestingly in this study, the oocytes of females in the high-quality group had on average smaller lipid
495 droplets with less size variation compared to those with low-quality group. Especially at DHP and
496 ovulation the difference between the groups was significant. The observed variation in lipid droplet
497 size, in conjunction with the hormone receptor expression pattern in the same females, suggests that
498 treatment administrated at an earlier stage enhanced maturational competence and the subsequent
499 developmental capacity of the egg/embryo. In general, limited information is available on the
500 mechanisms of oocyte lipid coalescence as well as on the role of this physiologically important process
501 in the subsequent survival and development of the eggs and embryos. Our findings, substantiating
502 insight in the underlying hormonal processes, suggest lipid droplet diameter to be an accurate,
503 quantitative indicator of maturation status in European eel.

504 In conclusion, ovarian *fshr*, *lhcr1*, *esr2a* and *gperb* were differentially expressed across sampling
505 points between females presenting high and low egg quality. A mismatch between the timing of
506 hormone injections and the expression of some of these genes may therefore influence the follicle's
507 ability to respond to treatment. Furthermore, changes in hormone receptor expression were associated
508 with changes in oocyte maturation status. Here, the average size of lipid droplets in advanced oocytes
509 was smaller in the high- compared to low-quality group throughout the maturation process, indicating
510 that the timing of hormonal treatments can be optimized, thereby optimizing procedures in assisted
511 reproduction of the endangered European eel.

512 **Acknowledgements**

513 We thank P. Lauesen, Billund Aquaculture Service, C. Graver, Danish Aquaculture Organisation, and
514 M. Krüger-Johnsen, Technical University of Denmark, for their assistance conducting experiments and
515 sampling, H. Johnsen, Nofima, for laboratory assistance, and A. Nielsen for guidance on statistical
516 methods. This work was supported by the European Commission's 7th Framework Programme (Grant
517 number 245257: *Reproduction of European Eel: Towards a self-sustained Aquaculture*, PRO-EEL) and
518 Innovation Fund Denmark (Grant number 11-2013-3: *Eel Hatchery Technology for a Sustainable*
519 *Aquaculture* (EEL-HATCH)), and a travel grant from the European Cooperation in Science and
520 Technology (COST, Grant number FA1205: AQUAGAMETE) to the first author in order to perform
521 gene expression studies at BOREA, France.

522 **References**

- 523 Andersson, E., Nijenhuis, W., Male, R., Swanson, P., Bogerd, J., Taranger, G.L., Schulz, R.W., 2009.
524 Pharmacological characterization, localization and quantification of expression of gonadotropin
525 receptors in Atlantic salmon (*Salmo salar* L.) ovaries. *Gen. Comp. Endocrinol.* 163, 329-339.
- 526 Asturiano, J.F., Pérez, L., Garzón, D.L., Marco-Jiménez, F., Peñaranda, D.S., Vicente, J.S., Jover, M.,
527 2004. Physio-chemical characteristics of seminal and development of media and methods for the
528 cryopreservation of European eel sperm. *Fish. Physiol. Biochem.* 30, 283-293.
- 529 Bobe, J., Maugars, G., Nguyen, T., Rime, H., Jalabert, B., 2003. Rainbow trout follicular maturational
530 competence acquisition is associated with an increased expression of follicle stimulating hormone
531 receptor and insulin-like growth factor 2 messenger RNAs. *Mol. Reprod. Dev.* 66, 46-53.
- 532 Butts, I.A.E., Sørensen, S.R., Politis, S.N., Pitcher, T.E., Tomkiewicz, J., 2014. Standardization of
533 fertilization protocols for the European eel, *Anguilla anguilla*. *Aquaculture* 426-427, 9-13.
- 534 Butts, I.A.E., Sørensen, S.R., Politis, S.N., Tomkiewicz, J., 2016. First-feeding by European eel larvae:
535 A step towards closing the life cycle in captivity. *Aquaculture* 464, 451-458.

536 Chauvigné, F., Zapater, C., Gasol, J.M., Cerdà, J., 2014. Germ-line activation of the luteinizing
537 hormone receptor directly drives spermiogenesis in a nonmammalian vertebrate. Proc. Natl. Acad.
538 Sci. U.S.A. 111, 1427-1432.

539 Conti, M., Andersen, C.B., Richard, F., Mehats, C., Chun, S.Y., Horner, K., Jin, C., Tsafirri, A., 2002.
540 Role of cyclic nucleotide signaling in oocyte maturation. Mol. Cell. Endocrinol. 187, 153-159.

541 da Silva, F.F.G., Støttrup, J.G., Kjørsvik, E., Tveiten, H., Tomkiewicz, J., 2016. Interactive effects of
542 dietary composition and hormonal treatment on reproductive development of cultured females
543 eel, *Anguilla anguilla*. Anim. Reprod. Sci. 171, 17-26.

544 Dufour S, Delerue-Le Belle N, Baloché, S., Fontaine Y.A., 1989. Positive feedback control by the
545 gonads on gonadotropin (GTH) and gonadoliberin (GnRH) levels in experimentally matured
546 female silver eels, *Anguilla anguilla*. Fish. Physio. Biochem. 7, 157-162.

547 Dufour S, Delerue-Le Belle N, Fontaine Y.A., 1983. Development of a heterologous
548 radioimmunoassay for eel (*Anguilla anguilla*) gonadotropin. Gen. Comp. Endocrinol. 49, 404-
549 413.

550 Dufour S., Lopez E., Le Menn F., Delerue-Le Belle N., Baloché S., Fontaine Y.A., 1988. Stimulation
551 of gonadotropin release and ovarian development, by the administration of a gonadoliberin
552 agonist and of dopamine antagonists, in female silver eel pretreated with estradiol. Gen. Comp.
553 Endocrinol. 70, 20-30.

554 Falkenstein, E., Tillmann, H.C., Christ, M., Feuring, M., Wehling, M., 2000. Multiple actions of steroid
555 hormones - a focus on rapid, nongenomic effects. Pharmacol. Rev. 52, 513-556.

556 Finet, B., Jalabert, B. and Garg, S.K., 1988. Effect of defolliculation and 17 α -hydroxy, 20 β -
557 dihydroprogesterone on cyclic AMP level in full-grown oocytes of the rainbow trout, *Salmo*
558 *gairdneri*. Gamete Res. 19, 241-252.

559 Fontaine, M., Bertrand, E., Lopez, E., Callamand, O., 1964 Sur la maturation des organes Génitaux de
560 l'Anguille femelle (*Anguilla anguilla* L.) et l'émission spontanée des œufs en aquarium. C. R.
561 Acad. Sci. 259, 2907-2910.

562 Hawkins, M.B., Thornton, J.W., Crews, D., Skipper, J.K., Dotte, A., Thomas, P., 2000. Identification
563 of a third distinct estrogen receptor and reclassification of estrogen receptors in teleosts. Proc.
564 Natl. Acad. Sci. USA 97, 10751-10756.

- 565 Ijiri, S., Kazeto, Y., Takeda, N., Chiba, H., Adachi, S., Yamauchi, K., 1995. Changes in serum steroid
566 hormones and steroidogenic ability of ovarian follicles during artificial maturation of cultivated
567 Japanese eel *Anguilla japonica*. *Aquaculture* 135, 3-16.
- 568 Jalabert, B., Finet, B., 1986. Regulation of oocyte maturation in the rainbow trout, *Salmo gairdneri*:
569 role of cyclic AMP in the mechanism of action of the maturation inducing steroid (MIS), 17 α
570 hydroxy, 20 β -dihydroprogesterone. *Fish. Physiol. Biochem.* 2, 65-74.
- 571 Jeng, S.R., Yueh, W.S., Chen, G.R., Lee, Y.H., Dufour, S., Chang, C.F., 2007. Differential expression
572 and regulation of gonadotropins and their receptors in the Japanese eel, *Anguilla japonica*. *Gen.*
573 *Comp. Endocrinol.* 154, 161-173.
- 574 Kagawa, H., Sakurai, Y., Horiuchi, R., Kazeto, Y., Gen, K., Imaizumi, H., Masuda, Y., 2013.
575 Mechanism of oocyte maturation and ovulation and its application to seed production in the
576 Japanese eel. *Fish Physiol. Biochem.* 39, 13-17.
- 577 Kagawa, H., Tanaka, H., Ohta, H., Unuma, T., Nomura, K., 2005. The first success of glass eel
578 production in the world: basic biology on fish reproduction advances new applied technology in
579 aquaculture. *Fish Physiol. Biochem.* 31, 193-199.
- 580 Kazeto Y, Kohara M, Tosaka R, Gen K, Yokoyama M, Miura C, Miura, T., Adachi, S., Yamauchi, K.,
581 2012. Molecular characterization and gene expression of Japanese eel (*Anguilla japonica*)
582 gonadotropin receptors. *Zool. Sci.* 29, 204-211.
- 583 Kobayashi, T., Pakarinen, P., Torgersen, J., Huhtaniemi, I., Andersen, Ø., 2008. The gonadotropin
584 receptors FSHR and LH-R of Atlantic halibut (*Hippoglossus hippoglossus*) -2. Differential
585 follicle expression and asynchronous oogenesis. *Gen. Comp. Endocrinol.* 156, 595-602.
- 586 Kwok, H.F., So, W.K., Wang, Y., Ge, W., 2005. Zebrafish gonadotropins and their receptors: I.
587 Cloning and characterization of zebrafish follicle-stimulating hormone and luteinizing hormone
588 receptors- evidence for their distinct functions in follicle development. *Biol. Reprod.* 72, 1370-
589 1381.
- 590 Lafont, A.-G., Rousseau, K., Tomkiewicz., Dufour, S., 2016. Three nuclear and two membrane
591 estrogen receptors in basal teleosts, *Anguilla* sp.: Identification, evolutionary history and
592 differential expression regulation. *Gen. Comp. Endocrinol.* 235, 177-191.

593 Lokman, P.M., Detlef, H.R., Davie, P.S., Young, G., 2003. The physiology of silvering in Anguillid
594 eels: androgens and control of metamorphosis from the yellow to silver stage. In: K. Aida, K.
595 Tsukamoto & K. Yamauchi (Eds.), *Eel Biology*, pp. 331-349. Springer Verlag, Tokyo.

596 Lubzens, E., Bobe, J., Young, G., Sullivan, C.V., 2017. Maternal investment in fish oocytes and eggs:
597 The molecular cargo and its contributions to fertility and early development. *Aquaculture* 472,
598 107-143.

599 Lubzens, E., Young, G., Bobe, J., Cerdà, J., 2010. Oogenesis in teleost: How fish eggs are formed.
600 *Gen. Comp. Endocrinol.* 165, 367-389.

601 Ma, C.H., Dong, K.W., Yu, K.L., 2000. cDNA cloning and expression of a novel estrogen receptor β -
602 subtype in goldfish (*Carassius auratus*). *Biochim. Biophys. Acta* 1490, 145-152.

603 Majumder, S., Das, S., Moulik, S.R., Mallick, B., Pal, P., Mukherjee, D., 2015. G-protein coupled
604 estrogen receptor (GPER) inhibits final oocyte maturation in common carp, *Cyprinus carpio*.
605 *Gen. Comp. Endocrinol.* 211, 28-38.

606 Maugars, G., Dufour, S., 2015. Demonstration of the Coexistence of Duplicated LH Receptors in
607 Teleosts, and Their Origin in Ancestral Actinopterygians. *Plos one* 10(8), e0135184.

608 Maugars, G., Schmitz, M., 2006. Molecular cloning and characterization of FSH and LH receptors in
609 Atlantic salmon (*Salmo salar* L.). *Gen. Comp. Endocrinol.* 149, 108-117.

610 Menuet, A., Pellegrini, E., Anglade, I., Blaise, O., Laudet, V., Kah, O., Pakdel, F., 2002. Molecular
611 characterization of three estrogen receptor forms in Zebrafish: binding characteristics,
612 transactivation properties, and tissue distributions. *Biol. Reprod.* 66, 1881-1892.

613 Morini, M., Peñaranda, D.D., Vílchez, M.C., Tveiten, H., Lafont, A.G., Dufour, S., Pérez, L.,
614 Asturiano, J.F., 2017. The expression of nuclear and membrane estrogen receptors in the
615 European eel throughout spermatogenesis. *Comp. Biochem. Physiol. A* 203, 91-99.

616 Mosselman, S., Polman, J., Dijkema, R., 1996. ER beta: identification and characterization of a novel
617 human estrogen receptor. *FEBS Lett.* 392, 49-53.

618 Mylonas, C.C., Fostier, A., Zanuy, S., 2010. Broodstock management and hormonal manipulations of
619 fish reproduction. *Gen. Comp. Endocrinol.* 165, 516-534.

620 Mylonas, C.C., Zohar, Y., 2007. Promoting oocyte maturation, ovulation and spawning in farmed fish.
621 In: Babin, P.J., Cerdà, J., Lubzens, E. (Eds.), *The fish oocyte: from basic studies to*
622 *biotechnological applications*. Springer, Dordrecht, The Netherlands. pp. 437-474.

623 Nagae, M., Todo, T., Gen, K., Kato, Y., Young, G., Adachi, S., Yamauchi, K., 1996. Molecular cloning
624 of the cDNAs encoding pituitary glycoprotein hormone alpha- and gonadotropin II beta-subunits
625 of the Japanese eel, *Anguilla japonica*, and increase in their mRNAs during ovarian development
626 induced by injection of chum salmon pituitary homogenate. *J. Mol. Endocrinol.* 16, 171-181.

627 Nagler, J.J., Cavileer, T., Sullivan, J., Cyr, D.G., Rexroad, C., 2007. The complete nuclear estrogen
628 receptor family in the rainbow trout: discovery of the novel ER α 2 and both ER β isoforms. *Gene*
629 392, 164-173.

630 Nelson, E.R., Habibi, H.R., 2010. Functional significance of nuclear estrogen receptor subtypes in the
631 liver of goldfish. *Endocrinology* 151, 1668-1676.

632 Norman, A.W., Mizwicki, M.T., Norman, D.P., 2004. Steroid-hormone rapid actions, membrane
633 receptors and a conformational ensemble model. *Nat. Rev. Drug Discovery* 3, 27-41.

634 Nyuji, M., Kitano, H., Shimizu, A., Lee, J.M., Kusakabe, T., Yamaguchi, A., Matsuyama, M., 2013.
635 Matsuyama M. Characterization, localization, and stage-dependent gene expression of
636 gonadotropin receptors in chub mackerel (*Scomber japonicus*) ovarian follicles. *Biol Reprod.* 88,
637 148.

638 Ohta, H., Kagawa, H., Tanaka, H., Okuzawa, K., Hirose, K., 1996. Changes in fertilization and
639 hatching rates with time after ovulation induced by 17, 20 β -dihydroxy-4-pregnen-3-one in the
640 Japanese eel, *Anguilla japonica*. *Aquaculture* 139, 291-301.

641 Ohta, H., Kagawa, H., Tanaka, H., Okuzawa, K., Linuma, N., Hirose, K., 1997. Artificial induction of
642 maturation and fertilization in the Japanese eel, *Anguilla japonica*. *Fish Physiol. Biochem.* 17,
643 163-169.

644 Palstra, A.P., Cohen, E.G.H., Niemantsverdriet, P.R.W., van Ginneken, V.J.T., van den Thillart,
645 G.E.E.J.M., 2005. Artificial maturation and reproduction of European silver eel: development of
646 oocytes during final maturation. *Aquaculture* 249, 533-547.

647 Pang, Y., Dong, J., Thomas, P., 2008. Estrogen signaling characteristics of Atlantic croaker G protein-
648 coupled receptor 30 (GPR30) and evidence it is involved in maintenance of oocyte meiotic arrest.
649 *Endocrinol.* 149, 3410-3426.

650 Pang, Y., Thomas, P., 2009. Involvement of estradiol-17 β and its membrane receptor, G protein
651 coupled receptor 30 (GPR30) in regulation of oocyte maturation in zebrafish, *Danio rario*. *Gen.*
652 *Comp. Endocrinol.* 161, 58-61.

653 Patiño, R., Yoshizaki, G., Thomas, P., Kagawa, H., 2001. Gonadotropic control of ovarian follicle
654 maturation: the two-stage concept and its mechanisms. *Comp. Biochem. Physiol.* 129, 427-439.

655 Patsoula E., Loutradis D., Drakakis P., Kallianidis K., Bletsas R. Michalakis S., 2001. Expression of
656 mRNA for the LH and FSH receptors in mouse oocytes and preimplantation embryos.
657 *Reproduction.* 121, 455-461.

658 Pedersen, B.H., 2003. Induced sexual maturation of European eel *Anguilla anguilla* and fertilisation of
659 the eggs. *Aquaculture.* 224, 323-338.

660 Pedersen, B.H., 2004. Fertilization of eggs, rate of embryonic development and hatching following
661 induced maturation of the European eel *Anguilla anguilla*. *Aquaculture* 237, 461-473.

662 Peñaranda, D.S., Pérez, L., Gallego, V., Barrera, R., Jover, M., Asturiano, J.F., 2010. European Eel
663 Sperm Diluent for Short-term Storage. *Reprod. Domest. Anim.* 45, 407-415.

664 Peyton, C., Thomas, P., 2011. Involvement of epidermal growth factor receptor signaling in estrogen
665 inhibition of oocyte maturation mediated through the G protein-coupled estrogen receptor (Gper)
666 in zebrafish (*Danio rerio*), *Biol. Reprod.* 85, 42-50.

667 Pierce J, Parsons TF., 1981. Glycoprotein hormones: structure and function. *J. Annu. Rev. Biochem.*
668 50, 465-495.

669 Prat, F., Sumpter, J.P., Tyler, C.R., 1996. Validation of radioimmunoassays for two salmon
670 gonadotropins (GTH I and GTH II) and their plasma concentrations throughout the reproductive
671 cycle in male and female rainbow trout (*Oncorhynchus mykiss*). *Biol. Reprod.* 54, 1375-1382.

672 Revankar, C.M., Cimino, D.F., Sklar, L.A., Arterburn, J.B., Prossnitz, E.R., 2005. A transmembrane
673 intracellular estrogen receptor mediates rapid cell signaling. *Science* 307, 1625-1630.

674 Rosenfeld, H., Meiri, I., Elizur, A., 2007. Gonadotropic regulation of oocyte development. In: Babin,
675 P.J., Cerdà, J., Lubzens, E. (Eds.), *The Fish Oocyte: From Basic Studies to Biotechnological*
676 *Applications.* Springer, Dordrecht, The Netherlands, pp. 175-202.

677 Saito, K., Lokman, P.M., Young, G., Ozaki, Y., Matsubara, H., Okumura, H., Kazeto, Y., Yoshiura, Y.,
678 Aida, K., Adachi, S., Yamauchi, K., 2003. Follicle-stimulating hormone β , luteinizing hormone
679 β , and glycoprotein hormone α subunit mRNA levels in artificially maturing Japanese eel
680 *Anguilla japonica* and naturally maturing New Zealand long-finned eel *Anguilla dieffenbachia*.
681 *Fish. Sci.* 69, 146-153.

682 Salmon, C., Nunez-Rodriguez, J., Marchelidon, J., Le Menn, F., Fontaine, Y.A., 1988. Gonadotropin
683 binding sites in eel ovary: Autoradiographic visualization and new data on specificity. *Reprod.*
684 *Nutr. Dévelop.* 28, 1165-1175.

685 Sambroni, E., Le Gac, F., Breton, B., Lareyre, J.J., 2007. Functional specificity of the rainbow trout
686 (*Oncorhynchus mykiss*) gonadotropin receptors as assayed in a mammalian cell line. *J.*
687 *Endocrinol.* 195, 213-228.

688 Schmitz, M., Aroua, S., Vidal, B., Le Belle, N., Elie, P., Dufour, S., 2005. Differential regulation of
689 luteinizing hormone and follicle-stimulating hormone expression during ovarian development
690 and under sexual steroid feedback in the European eel. *Neuroendocrinology* 81, 107-119.

691 Sørensen, S.R., Gallego, V., Pérez, L., Butts, I.A.E., Tomkiewicz, J., Asturiano, J.F., 2013. Evaluation
692 of methods to determine sperm density for the European eel, *Anguilla anguilla*. *Reprod. Domest.*
693 *Anim.* 48, 936-944.

694 Sørensen, S.R., Skov, P.V., Lauesen, P., Tomkiewicz, J., Bossier, P., Schryver, D.S., 2014. Microbial
695 interference and potential control in culture of European eel (*Anguilla anguilla*) embryos and
696 larvae. *Aquaculture* 426, 1-8.

697 Sørensen, S.R., Tomkiewicz, J., Munk, P., Butts, I.A.E., Nielsen, A., Lauesen, P., Graver, C., 2016.
698 Ontogeny and growth of early life stages of captive-bred European eel. *Aquaculture* 456, 50-61.

699 Stoeckel, J.N., 2000. A method for viewing the germinal vesicle in oocytes of commercial catfishes.
700 *North Amer. Journ. Aqua.* 62, 240-247.

701 Tang, H., Liu, Y., Li, J., Li, G., Chen, Y., Yin, Y., Guo, Y., Cheng, C.H.K., Liu, X., Lin, H., 2017. LH
702 signaling induced *ptgs2a* expression is required for ovulation in zebrafish. *Mol. Cell. Endocrinol.*
703 447, 125-133.

704 Thomas, P., Pang, Y., Filardo, E.J., Dong, J., 2005. Identity of an estrogen membrane receptor coupled
705 to a G protein in human breast cancer cells, *Endocrinology* 146, 624-632.

706 Tomkiewicz, J., 2012. Reproduction of European Eel in Aquaculture (REEL): Consolidation and new
707 production methods. DTU Aqua Report No 249-2012. National Institute Aquatic Resources,
708 DTU, 47 p.

709 Tomkiewicz, J., Kofoed, T.M.N., Pedersen, J.S., 2011. Assessment of testes development during
710 induced spermatogenesis in European eel *Anguilla anguilla*. *Mar. Coast. Fish.* 3, 106-118.

711 Tyler, C.R., Pottinger, T.G., Coward, K., Prat, F., Beresford, N., Maddix, S., 1997. Salmonid follicle-
712 stimulating hormone (GtH I) mediates vitellogenic development of oocytes in the rainbow trout,
713 *Oncorhynchus mykiss*. Biol. Reprod. 57, 1238-1244.

714 Unuma, T., Hasegawa, N., Sawaguchi, S., Tanaka, T., Matsubara, T., Nomura, K., Tanaka, H., 2011.
715 Fusion of lipid droplets in Japanese eel oocytes: Stage classification and its use as a biomarker
716 for induction of final oocyte maturation and ovulation. Aquaculture 322-323, 142-148.

717 Vidal, B., Pasqualini, C., Le Belle, N., Holland, M.C.H., Sbaihi, M., Vernier, P., Zohar, Y., Dufour, S.,
718 2004. Dopamine inhibits luteinizing hormone synthesis and release in the juvenile European eel:
719 a neuroendocrine lock for the onset of puberty. Biol. Reprod. 71, 1491-1500.

720 Wallace, R.A., 1985. Vitellogenesis and oocyte growth in nonmammalian vertebrates. In: Browder,
721 L.W. (Ed.), Developmental Biology. Plenum Press; New York, pp. 127-177.

722 Wallace, R.A., Selman, K., 1981. Cellular and dynamic aspects of oocyte growth in teleosts. Am. Zool.
723 21, 325-343.

724 Watson, C.S., Gametchu, B., 1999. Membrane-initiated steroid actions and the proteins that mediate
725 them. Proc. Soc. Exp. Biol. Med. 220, 9-19.

726 Yamamoto, K., Yamauchi, K., 1974. Sexual maturation of Japanese eel and production of eel larvae in
727 the aquarium. Nature. 251, 220-222.

728 Yamauchi, K., 1990. Studies of gonadal steroids involved in final gonadal maturation in the Japanese
729 eel, *Anguilla japonica*, a review. Inte. Revue. Ges. Hydrobiol. 75, 859-860.

730 Yoshiura, Y., Suetake, H., Aida, K., 1999. duality of gonadotropin in a primitive teleost, Japanese eel
731 (*Anguilla japonica*). 114, 121-131.

732 Zeleznik, A.J., Midgley, J.R., Reichert, J.R., 1974. Granulosa cell maturation in the rat: increased
733 binding of human chorionic gonadotropin following treatment with follicle-stimulating hormone
734 in vivo. Endocrinol. 95, 818-825.

735 Zohar, Y., Muñoz-Cueto, J.A., Elizur, A., Kah, O., 2010. Neuroendocrinology of reproduction in
736 teleost fish. Gen. Comp. Endocrinol. 165, 438-455.

737

738 **Legends**

739 **Figure 1: Oocytes at each sampling point.**

740 Photomicrographs showing examples of oocytes at the most advanced stage of development from the
741 biopsy samples taken a) before SPE priming, b) before DHP injection and c) ovulated eggs cleared in
742 Serra's liquid. GV: germinal vesicle and LD: lipid droplet.

743 **Figure 2: Expression of gonadotropin receptors.**

744 Boxplots representing expression of gonadotropin receptors (*fshr*, *lhcg1* and *lhcg2*) in ovarian
745 biopsies taken at the time of administration of the SPE primer (SPE) and DHP injection (DHP) as well
746 as ovulated eggs (EGG) in both high- (n=5) and low-quality (n=5) groups. The bold line inside each
747 box represents the median, lower and upper sides of each box represent the lower and upper quartile
748 (25% and 75%) and whiskers extend to the most extreme data point, no more than 1.5 times the
749 interquartile distance. Data points outside the boxplot are classed as "outliers". Significant differences
750 between groups are identified with "*" when $p < 0.05$; "***" when $p < 0.01$ and "****" when $p < 0.001$.

751 **Figure 3: Expression of estrogen receptors.**

752 Boxplots representing expression of nuclear (*esr1*, *esr2a*) and membrane estrogen receptors (*gpera* and
753 *gperb*) in ovarian biopsies taken at the time of administration of the SPE primer (SPE) and DHP
754 injection (DHP) as well as in ovulated eggs (EGG) in both high- (n=5) and low-quality (n=5) groups.
755 The bold line inside each box represents the median, lower and upper sides of each box represent the
756 lower and upper quartile (25% and 75%) and whiskers extend to the most extreme data point, no more
757 than 1.5 times the interquartile distance. Significant differences between groups are identified with "*"
758 when $p < 0.05$ and "***" when $p < 0.01$.

759 **Figure 4:**

760 Oocytes in three different maturational stages from an ovarian biopsy taken before the SPE priming
761 injection. 1: small and opaque oocytes; 2: large oocytes with darker cytoplasm; 3: large oocytes with
762 transparent cytoplasm and migratory nucleus.

763

764 **Figure 5: Lipid droplet diameter.**

765 Boxplot representing the lipid droplet diameter at the time of administration of the SPE priming (SPE)
766 and DHP injection (DHP), as well as in ovulated eggs (EGG) in both high- (n=5) and low-quality (n=5)
767 groups. The bold line inside each box represents the median, the lower and upper sides of each box
768 represent the lower and upper quartile (25% and 75%) and whiskers extend to the most extreme data
769 point which is no more than 1.5 times the interquartile distance. Data points outside the boxplot are
770 classed as “outliers”. Significant differences between egg groups are identified with an “*” when $p <$
771 0.05.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Table 1 Quantitative PCR primer sequences for gonadotropin receptors (*fshr*, *lhgr1* and *lhgr2*), nuclear estrogen receptors (*esr1*, *esr2a*, *esr2b*), membrane estrogen receptors (*gpera* and *gperb*) and reference gene (*18s*).

Name	Sequence (5' - 3')	Orientation	Accession number	Amplicon size (bp)	Reference
<i>fshr</i>	CCTGGTCGAGATAACAATCACC	Forward	LN831181	148	Maugars and Dufour (2015)
	AATCTTGGAGAAATCAGGCAGT	Reverse			
<i>lhgr1</i>	GCGGAAACACAGGGAGAAC	Forward	LN831182	155	Maugars and Dufour (2015)
	GGTTGAGGTACTGGAAATCGAAG	Reverse			
<i>lhgr2</i>	GTTTCCTGACCTATCGGCTATT	Forward	LN831183	132	Maugars and Dufour (2015)
	GGTTGAGGTACTGGAAATCGAAG	Reverse			
<i>esr1</i>	GCCATCATACTGCTCAACTCC	Forward	CUH82767	76	Lafont et al., (2015)
	CCGTAAAGCTGTCGTTTCAGG	Reverse			
<i>esr2a</i>	TGTGTGCCTCAAAGCCATTA	Forward	CUH82768	169	Lafont et al., (2015)
	AGACTGCTGCTGAAAGGTCA	Reverse			
<i>esr2b</i>	TGCTGGAATGCTGCTGGT	Forward	CUH82769	123	Lafont et al., (2015)
	CCACACAGTTGCCCTCATC	Reverse			
<i>gpera</i>	CAACTTCAACCACCGGGAGA	Forward	CUH82770	170	Lafont et al., (2015)
	TGACCTGGAGGAAGAGGGACA	Reverse			
<i>gperb</i>	AACCTGAACCACACGGAAA	Forward	CUH82771	170	Lafont et al., (2015)
	TGACCTGGAAGAAGAGGGACA	Reverse			
<i>18s</i>	CTCAACACGGGAAACCTCAC	Forward	FM946070	118	
	AGACAAATCGCTCCACCAAC	Reverse			

Table 2 Information and group assignment about female eels and resulting egg batches used in the present study, including initial length, (L_i) and body weight (BW_i), volume of eggs stripped, floating fraction (%), fertilization success (%), hatching success (%) and larval survival (%) at 3 days post hatch (dph).

Female No.	L_i (cm)	BW_i (g)	Eggs (g)	Floating	Fertilization	Hatching	Survival 3 dph	Group
1.	70	693	322	99	68	65	67	high-quality
2.	67	574	324	99	99	80	93	high-quality
3.	60	433	229	99	69	55	83	high-quality
4.	78	1014	335	95	97	19	13	high-quality
5.	86	1444	571	50	80	67	78	high-quality
6.	79	1019	372	25	84	4	0	low-quality
7.	63	414	200	90	15	5	0	low-quality
8.	92	1334	586	99	20	0	0	low-quality
9.	62	426	195	50	25	0	0	low-quality
10.	60	459	133	100	96	4	0	low-quality

Table 3 Pearson's correlation coefficient R and *p*-value for the linear regressions between gene expression at the time of SPE priming (SPE), DHP injection (DHP) and ovulated eggs (EGG), and hatching and fertilization success (n=10).

Gene	Time point	Hatching		Fertilization	
		R	<i>p</i>	R	<i>p</i>
<i>fshr</i>					
	SPE	0.23	0.529	-0.20	0.577
	DHP	-0.79	0.012	-0.46	0.213
	EGG	NA	NA	NA	NA
<i>lhcr1</i>					
	SPE	0.68	0.032	0.10	0.780
	DHP	0.36	0.314	-0.38	0.281
	EGG	-0.17	0.637	-0.17	0.636
<i>lhcr2</i>					
	SPE	0.00	0.999	0.42	0.223
	DHP	-0.24	0.512	0.40	0.250
	EGG	0.05	0.892	0.32	0.397
<i>esr1</i>					
	SPE	0.54	0.111	-0.11	0.762
	DHP	0.41	0.236	-0.33	0.351
	EGG	NA	NA	NA	NA
<i>esr2a</i>					
	SPE	0.60	0.067	0.05	0.894
	DHP	0.49	0.154	-0.11	0.762
	EGG	0.60	0.064	0.35	0.316
<i>gpera</i>					
	SPE	0.34	0.331	0.56	0.094
	DHP	0.29	0.419	0.46	0.183
	EGG	0.06	0.865	0.38	0.275
<i>gperb</i>					
	SPE	-0.44	0.204	-0.62	0.058
	DHP	-0.31	0.413	-0.21	0.584
	EGG	NA	NA	NA	NA